

Non-factorized calculation of the process ${}^3\text{He}(e, e'p)pn$ at medium energies

M. Alvioli

104 Davey Lab, The Pennsylvania State University, University Park, PA 16803, USA

C. Ciofi degli Atti and L.P. Kaptari*

Department of Physics, University of Perugia and Istituto Nazionale di Fisica Nucleare,
Sezione di Perugia, Via A. Pascoli, I-06123, Italy

(Dated: April 26, 2009)

The cross section and the transverse-longitudinal asymmetry A_{TL} of the three-body-break-up (3BBU) process ${}^3\text{He}(e, e'p)pn$ have been calculated by a non-factorized and parameter-free approach based upon realistic few-body wave functions corresponding to the AV18 interaction, treating the rescattering of the struck nucleon within a generalized eikonal approximation. The results of calculations exhibit a good agreement with recent JLab experimental data and show that the Final State Interaction (FSI) of the struck proton in the region of missing momentum $300 \lesssim p_m \lesssim 600 \text{ MeV}/c$ and removal energy $E_m \gtrsim p_m^2/4m_N$ is dominated by single nucleon rescattering, in agreement with nucleon-nucleon correlation predictions.

PACS numbers: 24.10.-i, 25.10.+s, 25.30.Dh, 25.30.Fj

In a previous paper [1] the cross section of the 2-body break up (2BBU) ${}^3\text{He}(e, e'p){}^2\text{H}$ recently measured at JLab [2] in the missing momentum range $0 \lesssim p_m \lesssim 1000 \text{ MeV}/c$, has been theoretically described by a fully non-factorized and parameter-free theoretical approach in which: i) initial state correlations (ISC) in the target nucleus ${}^3\text{He}$ have been taken care of by using state-of-the-art few-body wave functions obtained [3] by a variational solution of the Schrödinger equation containing realistic nucleon-nucleon interactions [4]; ii) FSI have been treated by a Generalized Eikonal Approximation (GEA) [5], which represents an extended Glauber approach (GA) based upon the evaluation of the relevant Feynman diagrams, which describe the rescattering of the struck nucleon in the final state, in analogy with the Feynman diagrammatic approach developed for the treatment of elastic hadron-nucleus scattering [6, 7]. The results obtained in [1], in agreement with other types of realistic calculations [8], have shown that: i) the FSI of the struck proton in the 2BBU channel can satisfactorily be described within single and double rescattering of the struck proton (cf. Fig. 1 of [1]); ii) the low momentum part ($p_m \lesssim 300 \text{ MeV}/c$) of the cross section is practically not affected by FSI and can even be described within a Factorized Approximation (FA), which assumes a factorization of the electron-Nucleus cross section into a product of the electron-nucleon cross section

and the nucleon Spectral Function; on the contrary at $300 \lesssim p_m \lesssim 1000 \text{ MeV}/c$, FSI becomes very important, particularly at "negative" values of the missing momentum (when ϕ , the angle between the scattering and reaction planes, equals zero), and the FA breaks down (cf. Fig. 2 of [1]); iii) in the region $300 \lesssim p_m \lesssim 700 \text{ MeV}/c$ the effects of FSI are almost entirely exhausted by single rescattering (we will come to this point later on) (cf. Fig. 2 of [1]); iv) the A_{TL} asymmetry is extremely sensitive to FSI and cannot be explained within a FA (cf. Fig. 3 of [1]). Following [1], we have undertaken the calculation of the 3-Body-Break-Up (3BBU) channel ${}^3\text{He}(e, e'p)pn$ using the same approach of [1]; our results will be compared with the experimental data of Ref. [9]. By considering the general case of a target nucleus A , the relevant quantities which characterize the process $A(e, e'p)(A-1)$ are the energy and momentum transfer ν and Q^2 , respectively, the missing momentum $\mathbf{p}_m = \mathbf{q} - \mathbf{p}_1$ (i.e. the momentum of the recoiling system $A-1$), and the *missing energy* $E_m = \sqrt{P_{A-1}^2 + m_N^2} - M_A = \nu - T_{A-1} - T_{\mathbf{p}_1} = |E_A| - |E_{A-1}| + E_{A-1}^* = E_{min} + E_{A-1}^*$. Here, $\mathbf{p}_1$ is the momentum of the detected proton, $E_A(E_{A-1})$ the (negative) ground state energy of the target (recoiling) nucleus and E_{A-1}^* the intrinsic excitation energy of the latter ($E_{A-1}^* = E_m - |E_{min}|$). The cross section of the process has form

$$\frac{d^6\sigma}{d\Omega' dE' d\Omega_{p_1} dE_{p_1}} = \left| \frac{\mathbf{p}_1^2}{\frac{|\mathbf{p}_1|}{E_{p_1}} + \frac{|\mathbf{p}_1| - |\mathbf{q}| \cos \theta}{E_{A-1}^*}} \right| \frac{dE_m}{dE_{p_1}} \sigma_{Mott} \sum_i V_i W_i^A(\nu, Q^2, \mathbf{p}_m, E_m) \quad (1)$$

where $i \equiv \{L, T, TL, TT\}$, V_i are kinematical factors and

the nuclear structure functions W_i^A result from proper

combinations of the polarization vector of the virtual photon, ε_λ^μ , and the hadronic tensor, $W_{\mu\nu}^A$, the latter depending upon the nuclear current operators $\hat{J}_\mu^A(0)$. We consider the interaction of the incoming virtual photon γ^* with a bound nucleon (the active nucleon) of low virtuality ($p^2 \sim m_N^2$) in the quasi-elastic kinematics i.e. corresponding to $x \equiv Q^2/2m_N\nu \sim 1$. In quasi-elastic kinematics, the virtuality of the struck nucleon after γ^* -

absorption is also rather low and, provided $\mathbf{p}_1$ is sufficiently high, nucleon rescattering with the "spectator" $A-1$ can be described to a large extent in terms of multiple elastic scattering processes in the eikonal approximation. Let us now consider the process ${}^3\text{He}(e, e')pn$. In co-ordinate representation the initial three-body wave function is $\Phi_{1/2, \mathcal{M}_3}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3)$, whereas the wave function of the final state is

$$\Psi_f^* = \hat{\mathcal{A}} \left[\chi_{T_f, \tau_f} S_\Delta^{FSI}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3) e^{-i\mathbf{p}_1 \mathbf{r}_1} e^{-i\mathbf{P}_m(\mathbf{r}_2 + \mathbf{r}_3)/2} \Phi_{S_f \sigma_f}^{\mathbf{k}_{rel}^*}(\mathbf{r}_2 - \mathbf{r}_3) \chi_{\frac{1}{2}, \lambda_f}^+ \right] \quad (2)$$

where $\hat{\mathcal{A}}$ is a proper antisymmetrization operator, χ_{T_f, τ_f} denotes the isospin wave function, $\Phi_{S_f \sigma_f}^{\mathbf{k}_{rel}^*}(\mathbf{r}_2 - \mathbf{r}_3)$ describes the relative motion of two interacting particles in

the continuum and, eventually, $\chi_{\frac{1}{2}, \lambda_f}^+$ is the spin wave function of the struck nucleon. The quantity

$$S_\Delta^{FSI}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3) = 1 + T_{(1)}^{FSI}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3) + T_{(2)}^{FSI}(\mathbf{r}_1, \mathbf{r}_2, \mathbf{r}_3) \quad (3)$$

is the GEA S-matrix, which describes both the case of no FSI, as well as single and double rescattering of the struck nucleon off the spectator nucleons [5, 10]. By introducing the corresponding Jacobi coordinates $\mathbf{R}$, $\boldsymbol{\rho}$ and $\mathbf{r}$, the relevant matrix elements, can be computed solely in terms of relative coordinates $\boldsymbol{\rho}$ and $\mathbf{r}$. For ease of presentation we will give explicit expressions of the single rescattering contribution only, which has the form $T_{(1)}^{FSI}(\boldsymbol{\rho}, \mathbf{r}) = - \sum_{i=2}^3 \theta(z_i - z_1) e^{i\Delta_z(z_i - z_1)} \Gamma(\mathbf{b}_1 - \mathbf{b}_i)$, where

$\mathbf{r}_i \equiv (z_i, \mathbf{b}_i)$. It can be seen that, unlike the usual GA, the GEA S-matrix gets also a contribution from a parallel momentum Δ_z of pure nuclear origin depending upon the external kinematics and the removal energy of the struck proton (within the "frozen approximation" $\Delta_z = 0$, and the usual Glauber profile is recovered). By assuming that the nuclear current operator is a sum of nucleonic currents $\hat{j}_\mu(i)$, its momentum space representation resulting from Feynman diagrams can be written as follows

$$\begin{aligned} J_\mu^3 &= \sum_\lambda \int \frac{d\mathbf{p}}{(2\pi)^3} \frac{d\mathbf{k}}{(2\pi)^3} S_\Delta^{FSI}(\mathbf{p}, \mathbf{k}) \langle \lambda_f | j_\mu(\mathbf{k} - \mathbf{p}_m; \mathbf{q}) | \lambda \rangle \mathcal{O}(\mathbf{p}_m - \mathbf{k}, \mathbf{p}, \mathbf{k}_{rel}; \mathcal{M}_3, S_f, \sigma_f, \lambda) = \\ &= J_\mu^{3(PWIA)} + J_\mu^{3(1)} + J_\mu^{3(2)}, \end{aligned} \quad (4)$$

where $S_\Delta^{FSI}(\mathbf{p}, \mathbf{k}) = \int d\mathbf{r} d\boldsymbol{\rho} e^{-i\mathbf{p}\mathbf{r}} e^{i\mathbf{k}\boldsymbol{\rho}} S_\Delta^{FSI}(\boldsymbol{\rho}, \mathbf{r})$, is the Fourier transform of the GEA S-matrix (Eq. (3)), $\mathcal{O}$ the

nuclear overlap in momentum space

$$\mathcal{O}(\mathbf{p}_m - \mathbf{k}, \mathbf{p}, \mathbf{k}_{rel}; \mathcal{M}_3, S_f, \sigma_f, \lambda) = \int d\mathbf{p} d\mathbf{r} e^{i(\mathbf{p}_m - \mathbf{k})\mathbf{r}} e^{i\mathbf{p}\mathbf{r}} \Phi_{\frac{1}{2}, \mathcal{M}_3}(\boldsymbol{\rho}, \mathbf{r}) \Phi_{S_f, \sigma_f}^{\mathbf{k}_{rel}^*}(\mathbf{r}) \chi_{\frac{1}{2}, \lambda}^+ \quad (5)$$

and, eventually, $\langle \lambda_f | j_\mu(\mathbf{k} - \mathbf{p}_m; \mathbf{q}) | \lambda \rangle$ is the nucleon cur-

rent operator, with λ and λ_f being the spin projec-

tions of the struck proton before and after γ^* absorption. Note, that in eq. (5) the quantity $\mathbf{\kappa} - \mathbf{p}_m \equiv \mathbf{k}_1$ represents exactly the momentum of the struck proton before the electromagnetic interaction, with $\mathbf{\kappa}$ being the momentum transfer in single rescattering. In absence of any FSI the momentum space S-matrix is $S_{\Delta}^{FSI}(\mathbf{\kappa}, \mathbf{p}) = (2\pi)^6 \delta(\mathbf{\kappa}) \delta(\mathbf{p})$ and only the PWIA con-

tribution $J_{\mu}^{3(PWIA)}$ survives in Eq. (4). When FSI is active, the contributions from single and double rescattering have to be taken into account; let us consider the former: it results from the single scattering momentum space term of $S_{\Delta}^{FSI}(\mathbf{p}, \mathbf{\kappa})$, which has the following form

$$T_{(1)}^{FSI}(\mathbf{p}, \mathbf{\kappa}) = \frac{(2\pi)^4}{k^*} \frac{f_{NN}(\mathbf{\kappa}_{\perp})}{\mathbf{\kappa}_{\parallel} + \Delta_z - i\varepsilon} \left[\delta\left(\mathbf{p} - \frac{\mathbf{\kappa}}{2}\right) + \delta\left(\mathbf{p} + \frac{\mathbf{\kappa}}{2}\right) \right] \quad (6)$$

where $f_{NN}(\mathbf{\kappa}_{\perp})$ (the Fourier transform of the profile function $\Gamma(\mathbf{b})$) represents the elastic scattering amplitude of two nucleons with center-of-mass momentum k^* . Placing Eq. (6) in Eq. (4) the single scattering contribution $J_{\mu}^{3(1)}$ to the nuclear current is obtained as follows


$$J_{\mu}^{3(1)} = \sum_{\lambda} \int \frac{d\mathbf{\kappa}}{(2\pi)^2 k^*} \langle \lambda_f | j_{\mu}(\mathbf{k}_1; \mathbf{q}) | \lambda \rangle \frac{f_{NN}(\mathbf{\kappa}_{\perp})}{\mathbf{\kappa}_{\parallel} + \Delta_{\parallel} - i\varepsilon} \times \\ \left[\mathcal{O}(-\mathbf{k}_1, \frac{\mathbf{\kappa}}{2}; \mathcal{M}_3, S_f, \sigma_f, \lambda) + \mathcal{O}(-\mathbf{k}_1, -\frac{\mathbf{\kappa}}{2}; \mathcal{M}_3, S_f, \sigma_f, \lambda) \right]; \quad (7)$$

where $\mathbf{k}_1$, as already mentioned, is the momentum of the proton before γ^* absorption and $\mathbf{\kappa} = \mathbf{k}_1 + \mathbf{q} - \mathbf{p}_1 = \mathbf{k}_1 + \mathbf{p}_m$ is the momentum transfer in the NN rescattering. The longitudinal part of the nucleon propagators has been computed using the relation $[\mathbf{\kappa}_{\parallel} + \Delta_z \pm i\varepsilon]^{-1} = \mp i\pi \delta(\mathbf{\kappa}_{\parallel} + \Delta_z) + PV[\mathbf{\kappa}_{\parallel} + \Delta_z]^{-1}$. Note that in the eikonal approximation the trajectory of the fast nucleon is a straight line so that all "longitudinal" and "perpendicular" components are defined with respect to this trajectory, which means that the z axis has to be directed along the momentum of the detected fast proton. Looking at the structure of Eq. (7), it can be seen that, due to the coupling of the nucleonic current operator $\langle \lambda_f | j_{\mu}(\mathbf{k}_1; \mathbf{q}) | \lambda \rangle$ with the nuclear overlap integral, a factorized form for Eq. (7) cannot be obtained; the same holds for the double-scattering contribution and, consequently, for the cross section. However, as shown in Ref. [10], if the longitudinal part of the nucleonic current can be disregarded, the factorization form can approximately be recovered. Using the above formalism, and including the contribution from double rescattering $J_{\mu}^{3(2)}$, the cross section (Eq. (1)) and the left-right asymmetry defined by

$$A_{TL} = \frac{d\sigma(\phi = 0^\circ) - d\sigma(\phi = 180^\circ)}{d\sigma(\phi = 0^\circ) + d\sigma(\phi = 180^\circ)}. \quad (8)$$

have been calculated. Following de Forest's prescription [11], the "CC1" form of the nucleon current operator has been adopted; the elastic amplitude f_{NN} has been chosen in the usual form $f_{NN}(\mathbf{\kappa}_{\perp}) = k^* \frac{\sigma^{tot}(i+\alpha)}{4\pi} e^{-b^2 \mathbf{\kappa}_{\perp}^2/2}$, where the slope parameter b , the total nucleon-nucleon cross section σ^{tot} and the ratio α of the real to the

imaginary parts of the forward scattering amplitude, were taken from world's experimental data. The results of our calculations are shown in Figs. 1, 2 and 3. In Fig. 1 the full non-factorized cross section at two values of p_m is compared with the PWIA result and with the experimental data; in this Figure the role played by single and double rescattering is clearly illustrated. In Fig. 2 the factorized and non-factorized results are compared at $p_m = 440 \text{ MeV}/c$ (similar results are obtained at higher values of p_m). Finally in Fig. 3 the asymmetry A_{TL} is exhibited. As in Ref. [1], no approximations have been made in the evaluation of the single and double rescattering contributions: intrinsic coordinates have been used and the energy dependence of the profile function has been taken into account in the properly chosen CM system of the interacting pair. Note that the experimental kinematics has been intentionally chosen so as to cover the theoretically predicted two nucleon correlation region characterized by a peak position given by $E_2^* = E_m - E_{min} \sim 2m_N \{ \sqrt{(1/2)[1 + \sqrt{1 + (p_m/m_N)^2}] - 1} \} \rightarrow p_m^2/4m_N$ in the non relativistic limit. It can be seen that the location of the experimental peaks agrees with such a prediction. The amplitude of the peak is largely affected by FSI, whose inclusion brings the theoretical calculation in very good agreement with the experimental data. The relevant effects of FSI can be explained by the fact that the kinematics of the experiment is such that the missing momentum is always almost perpendicular to the momentum of the final proton and FSI is maximized. Concerning the effects of the latter, an interesting ob-



A_{TL} of the three-body-break-up (3BBU) process ${}^3\text{He}(e, e'p)pn$ have been calculated by a non-factorized and parameter-free approach based upon realistic few-body wave functions corresponding to the $AV18$ interaction, treating the rescattering of the struck nucleon within a generalized eikonal approximation. The results of calculations exhibit a good agreement with recent JLab experimental data and show that the Final State Interaction (FSI) of the struck proton in the region of missing momentum $300 \lesssim p_m \lesssim 600 \text{ MeV}/c$ and removal energy $E_m \gtrsim p_m^2/4m_N$ is dominated by single nucleon rescattering, in agreement with nucleon-nucleon correlation predictions.

FIG. 1: The differential cross section (Eq. (1)) for the process ${}^3\text{He}(e, e'p)pn$ calculated at two values of the missing momentum *vs.* the excitation energy of the two-body final state $E_2^* = E_m - E_{min}$ ($E_{min} = 2m_p + m_n - M_3$). Dotted lines: PWIA approximation; dashed and solid lines: non-factorized calculations with single and double rescattering in the final state, respectively. Experimental data from [9].

servation is in order: it can be seen that in the region of the correlation peak, and at higher values of E_2^* , the effects from double rescattering is very tiny; the same appears to be the case in the 2BBU channel at $300 \lesssim p_m \lesssim 700 \text{ MeV}/c$. We consider this as another evidence that in these regions of missing momentum and energy FSI mainly occurs within the correlated pair. In Fig. 2 the factorized and non-factorized results are compared at $p_m = 440 \text{ MeV}/c$; it can be seen that the non-factorized calculation differs by 5-7% from the factorized one and better agrees with the experimental data. Eventually, in Fig. 3 the transverse-longitudinal asymmetry A_{TL} (Eq. (8)), calculated within the PWIA




FIG. 2: Comparison at $p_m = 440 \text{ MeV}$ of the results of calculations performed within the non-factorized approach (NFA) and the factorization approximation (FA).

and taking into account FSI, is presented. It is well known that when the explicit expressions of V_i and W_i^A are placed in Eq. (8), the numerator is proportional to the transverse-longitudinal response W_{TL} , whereas the denominator does not contain W_{TL} at all, which means that A_{TL} is a measure of the relevance of the transverse-longitudinal response relative to the other responses. In the eN cross section the behavior of the asymmetry A_{TL} is known to be a negative and decreasing function of the missing momentum [11] and the same behavior should be expected in eA scattering within the PWIA. The asymmetry presented in Fig. 3 clearly shows that a measurement of the 3BBU channel at $\phi = 0^\circ$ would provide results not very different from the ones obtained at $\phi = 180^\circ$. To sum up, we have calculated in momentum space the cross section of the processes ${}^3\text{He}(e, e'p)pn$, using realistic ground-state two- and three-body wave functions, treating the FSI of the struck nucleon with the spectator nucleon pair within the generalized eikonal approximation. As in the case of the 2BBU calculation [1], our approach is a parameter free one, for it only requires the knowledge of the nuclear wave functions, the FSI factor being fixed directly by NN scattering data. At the same time, calculations are very involved mainly because of the complex structure of the wave function of Ref. [3], which has to be firstly transformed to momentum space and then used in the calculations of multidimensional integrals, including also the computation of principal values. The main results of our calculations can be summarized as follows: i) a good agreement between theoretical calculations and experimental data on the 3BBU process ${}^3\text{He}(e, e'p)pn$ has been achieved; ii) FSI effects are very relevant, but in the region of the correlation peak ($E_2^* \simeq \frac{p_m^2}{4m}$) and at higher energies ($E_2^* \geq \frac{p_m^2}{4m}$), they can be accounted for by single rescattering, the double rescattering contribution




FIG. 3: The A_{TL} asymmetry (Eq. (8)) calculated in correspondence of two values of p_m vs. the excitation energy of the two-body final state.

being negligible; this is an implicit confirmation of the two-nucleon correlation picture, according to which the uncorrelated nucleon (the A-2 system) is far away from the correlated pair and practically does not contribute to the short-range rescattering processes in the final state; iii) factorized and non-factorized results may differ at

$\phi = 180^\circ$ by at most 10%.

The authors are indebted to the Pisa Group for making available the variational three-body wave functions. Thanks are due to S. Gilad, M. Sargsian, S. Scopetta, and M. Strikman for useful discussions. L.P.K. is indebted to the Italian Ministry of University and Research (MIUR) for a grant within the program "Rientro dei Cervelli". He thanks the University of Perugia and INFN, Sezione di Perugia, for warm hospitality. M.A. is supported by DOE grant under contract DE-FG02-93ER40771

* On leave from Bogoliubov Lab. Theor. Phys., 141980, JINR, Dubna, Russia

- [1] C.Ciofi degli Atti, L.P. Kaptari, Phys. Rev. Lett. **100** (2008) 122301.
- [2] M. M. Rvachev et al., Phys. Rev. Lett. **94** (2005) 192302.
- [3] A. Kievsky, S. Rosati and M. Viviani, Nucl. Phys. **A551** (1993) 241 and *Private communication*.
- [4] R. B. Wiringa, V. G. J. Stoks and R. Schiavilla, Phys. Rev. **C51** (1995) 38.
- [5] L.L. Frankfurt, M.M. Sargsian, M.I. Strikman, Phys. Rev. **C56** (1997) 1124; M.M. Sargsian, Int. J. Mod. Phys. **E10**(2001)405.
- [6] V.N.Gribov, Sov. Phys. JETP, **30** (1970) 709.
- [7] L. Bertocchi, Nuovo Cimento, **11A** (1972) 45.
- [8] R. Schiavilla, O. Benhar, A. Kievsky, L.E. Marcucci, M. Viviani, Phys. Rev. **C72** (2005) 064003.
- [9] F. Benmokhtar et al., Phys. Rev. Lett. **94** (2005) 082305.
- [10] C.Ciofi degli Atti, L.P. Kaptari, Phys. Rev. **C71** (2005) 024005; C.Ciofi degli Atti, L.P. Kaptari, Phys. Rev. Lett. **95** (2005) 052502.
- [11] T. de Forest Jr., Nucl. Phys. **A392** (1983) 232.